




4949 SW 41st Blvd.
Suites 40 & 50
Gainesville, FL 32608
Tel: (352) 372 7022
Fax: (352) 372 7066
admin@encorbio.com

Catalogue# CPCA-GAP43: Chicken polyclonal antibody to GAP43: GAP43

The Immunogen: GAP43 is very abundant nervous system protein which is found concentrated in neurons. One group discovered it as one of three proteins which becomes unregulated during the regeneration of the toad optic nerve (1). Three GAPs (Growth associated proteins) were discovered, and the number 43 comes from the apparent [SDS-PAGE](#) molecular weight of the one named GAP43. Later work showed that GAP43 does not run on SDS-PAGE in a fashion which accurately reflects its molecular weight, and that GAP43 proteins from different species may run at different apparent molecular weights. Partly due to these features GAP43 was independently discovered by several different groups and therefore has several alternate names, such as protein F1, pp46, neuromodulin, neural phosphoprotein B-50 and calmodulin-binding protein P-57. In each case the number reflects the apparent SDS-PAGE molecular weight, and underlines the unusual SDS-PAGE mobility properties of this molecule. Mammalian GAP43 proteins contains only 226-243 amino acids, and so the real molecular weight is 23.61-25.14KDa (to perform such calculations yourself see [this link](#)). GAP43 is one of many highly negatively charged extended molecules which lack well defined tertiary structure and contain few hydrophobic residues and which all run anomalously on SDS-PAGE. Other examples are CAP23, MARCKS, microtubule associated proteins MAP2 and tau and the neurofilament subunits. GAP43 has been extensively studied and is known to be a major protein kinase C substrate and to bind calmodulin avidly. GAP43 is anchored to the plasma membrane by palmitoylation modifications. The [HGNC](#) name for this protein is [GAP43](#).



Left: Western blots of extract of cow hippocampus stained with CPCA-GAP43. A prominent band running at ~43 kDa represents GAP43. **Right:** Mixed neuron-glia cultures stained with CPCA-GAP43 (green) and rabbit antibody to alpha-II spectrin (red), and DNA (blue). The GAP43 antibody stains numerous axonal and dendritic profiles in these cultures, clearly revealing the submembrane cytoskeleton and associated vesicles.

Antibody characteristics: The antibody was raised against the C-terminal peptide of rat and mouse GAP43, which is KEDPEADQEHA, to which an N-terminal Cysteine residue was added to allow chemical coupling to Keyhole Limpet Hemocyanin carrier protein. The corresponding C-terminal peptide in the human GAP43 sequence is KGEEREADQEHA, which is similar enough that our antibody binds to this also. The antibody is provided as an IgY preparation.

Suggestions for use: The IgY solution can be diluted to 1:500-1,000 for immunofluorescence staining and 1:1,000-5,000 for western blotting. On western blots look for a major band at about 43 kDa, though this may vary somewhat depending on the species and the acrylamide concentration of your SDS-PAGE system (see above). Store at 4°C or -20°C. Avoid repeat freezing and thawing.

References:

1. Skene JH, Willard M. Changes in axonally transported proteins during axon regeneration in toad retinal ganglion cells. [J Cell Biol. 89:86-95 \(1981\).](#)

Limitations: This product is for research use only and is not approved for use in humans or in clinical diagnosis.

[©EnCor Biotechnology Inc.](#) May 23, 2014.