

Ordering Information

Web www.encorbio.com
Email admin@encorbio.com
Phone 352-372-7022
Fax 352-372-7066

HGNC Name: VSNL1

UniProt: P62760

RRID: AB_2572399

Immunogen: Recombinant full length human protein

Format: Purified antibody at 1mg/mL in 50% PBS,
50% glycerol plus 5mM NaN₃

Storage: Store at 4°C for short term, for longer term
store at -20°C


Recommended dilutions:

WB: 500-1,000. IF/ICC and IHC: 1:500-1,000


References:

1. Hatakenaka S, Kuo CH, Miki N. Analysis of a distinctive protein in chick retina during development. *Brain Res.* 312:155-63 (1983).
2. Kuno T, et al. cDNA cloning of a neural visinin-like Ca(2+)-binding protein. *Biochem. Biophys. Res. Commun.* 184:1219-25 (1992).
3. Polymeropoulos MH, Ide S, Soares MB, Lennon GG. Sequence characterization and genetic mapping of the human VSNL1 gene, a homologue of the rat visinin-like peptide RNVP1. *Genomics* 29:273-5 (1995).
4. Bernstein HG, et al. Regional and cellular distribution of neural visinin-like protein immunoreactivities (VILIP-1 and VILIP-3) in human brain. *J. Neurocytol.* 28:655-62 (1999).
5. Paterlini M, Revilla V, Grant AL, Wisden W. Expression of the neuronal calcium sensor protein family in the rat brain. *Neuroscience* 99:205-16 (2000).
6. Laterza OF, et al. Identification of novel brain biomarkers. *Clin. Chem.* 9:1713-21 (2006).
7. Lee JM, et al. The brain injury biomarker VLP-1 is increased in the cerebrospinal fluid of Alzheimer disease patients. *Clin. Chem.* 10:1617-23 (2008).
8. Tarawneh R, et al. Visinin-like protein-1: diagnostic and prognostic biomarker in Alzheimer disease. *Ann Neurol.* 70:274-85 (2011).

Applications	Host	Isotype	Molecular Wt.	Species Cross-Reactivity
WB, IF/ICC, IHC	Mouse	IgG1	18kDa	Hu, Rt, Ms



Western blot analysis of different tissue lysates using mouse mAb to Visinin-like Protein 1 (VLP1), MCA-2D11, dilution 1:1,000 in green: [1] protein standard (red), [2] rat brain [3] rat cerebellum, and [4] mouse brain. The band at ~20kDa mark corresponds to the VLP1 protein.



Immunofluorescent analysis of rat cerebellum section stained with mouse mAb to VLP1, MCA-2D11, dilution 1:500, in red and costained with rabbit pAb to GFAP, [RPCA-GFAP](#), dilution 1:5,000 in green. Following transcardial perfusion of rat with 4% paraformaldehyde, brain was post fixed for 24 hours, cut to 45µm, and free-floating sections were stained with the above antibodies. The VLP1 antibody reveals protein expressed in granule cell membranes and in synapses in the white matter and molecular layers of the cerebellum. The GFAP antibody stains the processes of Bergmann glia and astroglia.

Background:

Visinin was originally isolated biochemically from chicken retina as a major protein of ~24kDa on SDS-PAGE gels (1). Following cloning and sequencing of visinin, several visinin like proteins were discovered by homology screening (2,3). One of these, [visinin-like protein 1](#) (VLP-1) is a low molecular weight protein which is very abundant in the nervous system and is found only in neurons, though different neurons have different levels of expression (4,5). The protein was discovered independently by several groups and is therefore also sometimes known as hippocalcin-like protein 3, HLP3, HPCAL3, HUVISL1, VILIP and VILIP-1. The protein belongs to the large superfamily of [calmodulin](#) and [parvalbumin](#) type proteins which function by binding Calcium ions. Calcium binding alters the conformation of these proteins and allow them to interact with other binding partners, the properties of which they may alter. Visinin-like protein 1 has four "EF hand" domains, which are negatively charged helix-turn-helix peptides which are responsible for Calcium binding. The protein is 191 amino acids in size and has a molecular weight on SDS-PAGE of 18kDa. The protein has recently been suggested to be a useful blood biomarker of Alzheimer's disease and traumatic brain injury (6-8).

The MCA-2D11 antibody was made against full length recombinant human VLP-1. It can be used to track this protein by ELISA, on western blots and in cells in culture and sections for IF, ICC and IHC. VLP-1 is heavily concentrated in cerebellar granule cells and in most other neuronal types. We manufacture an alternate mouse monoclonal antibody and also rabbit and chicken polyclonal antibodies to this protein, [MCA-3A9](#), [RPCA-VLP1](#) and [CPCA-VLP1](#) respectively.

FOR RESEARCH USE ONLY. NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE.

Abbreviation Key:

mAb—Monoclonal Antibody pAb—Polyclonal Antibody WB—Western Blot IF—Immunofluorescence ICC—Immunocytochemistry
IHC—Immunohistochemistry E—ELISA Hu—Human Mo—Monkey Do—Dog Rt—Rat Ms—Mouse Co—Cow Pi—Pig Ho—Horse Ch—Chicken
Dr—D. rerio Dm—D. melanogaster Sm—S. mutans Ce—C. elegans Sc—S. cerevisiae Sa—S. aureus Ec—E. coli.