C-FOS Biotechnology Inc.

MCA-2H2

Ordering Information Web www.encorbio.com Email admin@encorbio.com Phone 352-372-7022 Fax 352-372-7066

HGNC Name: FOS UniProt: P01100 RRID: AB_2571561 Immunogen: Full length recombinant human protein expressed in and purified from E. coli. Format: Purified antibody at 1mg/mL in 50% PBS, 50% glycerol plus 5mM NaN₃ Storage: Store at 4°C for short term, for longer term at -20°C. Recommended dilutions: WB: 1:500, IF/ICC or IHC: 1:500

References:

1. Mildle-Langosch K. The Fos family of transcription factors and their role in tumourigenesis. Eur. J. Cancer 41:2449-2461 (2005)..

2. Chiu R, et al. The c-Fos protein interacts with c-Jun/AP-1 to stimulate transcription of AP-1 responsive genes. Cell 54:541–52 (1988). 3. Karin M. The regulation of AP-1 activity by mitogen activated protein kinases. J Biol Chem. 270:16483-6 (1995).

4. Bossis G, et al. Down-regulation of c-Fos/c-Jun AP-1 dimer activity by sumoylation. Mol Cell Biol.25(16):6964-79 (2005).

5. Dragunow M, Faull R. The use of c-fos as a metabolic marker in neuronal pathway tracing. J. Neurosci. Mets. 29:261–265 (1989).

This is a new antibody but peer reviewed publications using it are beginning to come on line. So if you perform a Google Scholar search for "MCA-2H2" several publications will appear, or you can just select here.



Western blot analysis of cell lysates using mouse mAb to cFos, MCA-2H2, dilution 1:1,000, in green, and rabbit pAb to GAPDH, RPCA-GAPDH, dilution 1:20,000, in red, used as a loading control. [1] protein standard (red), [2] HeLa cells in serum free media. [3] HeLa cells stimulated with 20% fetal bovine serum for 2hrs after 36hrs in serum free media. [4] rat cortical neurons. [5] rat cortical neurons treated with membrane depolarization buffer for 5hrs. Multiple bands at 50-65kDa in stimulated or treated cell lysates correspond to different forms of the c-Fos proten. The single band at 37 kDa represents GAPDH protein.

Isotype Molecular Wt. Species Cross-Reactivity IgG1 heavy, к light 50-65kDa Hu, Rt, Ms



Immunofluorescent analysis of rat hippocampus section stained with mouse mAb to c-FOS, MCA-2H2, dilution 1:200, in red, and costained with rabbit pAb to FOX3/NeuN, RPCA-FOX3, dilution 1:3,000, in green. The blue is Hoechst staining of nuclear DNA. The MCA-2H2 antibody labels nuclei of spontaneously activated neurons, while FOX3/NeuN antibody stains nuclei and distal perikarya of most neurons.

Background:

The FOS gene and protein were originally identified as the transforming element in a viral oncogene. The transforming protein was named v-FOS, for viral FOS, and the normal cellular non-transforming proto-oncogene was called c-FOS, for cellular FOS. FOS is an acrynym for "FBJ murine osteogenic sarcoma", the virus in which the gene product was first discovered. The c-FOS protein is a normal gene acting as an on/off switch controlling the expression of many other genes. The v-FOS form is mutated to stay in the on position, this persistently activating other genes and promoting unregulated cell division. The unmutated c-FOS is an "immediate-early" gene, so-called because protein expression is usually very low but increases rapidly and transiently in response to a wide array of stimuli including serum, growth factors, tumor promoters, cytokines, and UV radiation. Newly expressed c-FOS protein associates with JUN family and other basic leucine-zipper (bZIP) proteins to create a variety of activator protein-1 (AP-1) complexes (1). AP-1 complexes specifically activate the expression of many other genes and so regulate cellular responses to stimuli which may result in cell proliferation, differentiation, neoplastic transformation, apoptosis, and response to stress (2). The regulated expression of c-FOS inhibits the AP-1 transcriptional activity (3,4). Since c-FOS expression is induced in neurons which are rapidly firing action potentials, appropriate c-Fos antibodies can be used to identify activated neurons in tissues (5). Using current techniques it is possible to follow processes of such cells or obtain data on their mRNA expression.

The MCA-2H2 antibody was made against recombinant full length human c-FOS expressed in and purified from *E. coli*. It can be used to identify activated cells in cell culture and in sections and to follow c-FOS expression in western blots of cell and tissue homogenates. The

antibody also works well on formalin fixed paraffin embedded sections, select the "Additional Info" for this data. The KD is 6.68 x 10.10 M,

Kon rate is $1.36 \times 10^{\circ}$ 1/MS and the Kdis rate is $9.12 \times 10^{\circ}$ 1/S, all indicative of unusually high affinity. The same recombinant immunogen was used to generate a rabbit polyclonal antibody to c-FOS, RPCA-c-Fos, which has similar properties.

FOR RESEARCH USE ONLY. NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE.

Abbreviation Key:

mAb—Monoclonal Antibody pAb—Polyclonal Antibody WB—Western Blot IF—Immunofluorescence ICC—Immunocytochemistry IHC—Immunohistochemistry E—ELISA Hu—Human Mo—Monkey Do—Dog Rt—Rat Ms—Mouse Co—Cow Pi—Pig Ho—Horse Ch—Chicken Dr—D. rerio Dm—D. melanogaster Sm—S. mutans Ce—C. elegans Sc—S. cerevisiae Sa—S. aureus Ec—E. coli.