


Ordering Information
 Web www.encorbio.com
 Email admin@encorbio.com
 Phone 352-372-7022
 Fax 352-372-7066

HGNC Name: MAP2
UniProt: P11137
RRID: AB_2737286
Immunogen: Recombinant human projection domain sequences, amino acids 377-1505, EnCor products [Prot-r-MAP2-P1](#), [Prot-r-MAP2-P2](#) and [Prot-r-MAP2-P3](#).
Format: Purified antibody at 1mg/mL in 50% PBS, 50% glycerol plus 5mM NaCl.
Storage: Store at 4°C for short term, for longer term at -20°C
Recommended dilutions:
 WB: 1:2,000. IF/ICC and IHC: 1:1,000-1:2,000.


References:

1. Dehmelt H, Halpain S. The MAP2/Tau family of microtubule-associated proteins. *Genome Biol.* 6:204 (2005).
2. Nunez J. Immature and mature variants of MAP2 and tau proteins and neuronal plasticity. *Trends Neurosci.* 11:477-9 (1998).
3. 2. Vallee R. A taxol-dependent procedure for the isolation of microtubules and microtubule-associated proteins (MAPs). *J. Cell Biol.* 92:435-42 (1992).
4. Goetz AK, et al. Temporally restricted substrate interactions direct fate and specification of neural precursors derived from embryonic stem cells. *PNAS* 103:11063-8 (2006).
5. Walton NM et al. Gliotypic neural stem cells transiently adopt tumorigenic properties during normal differentiation. *Stem Cells* 27:280-9 (2009).
6. Gasser A, et al. An ankyrinG-binding motif is necessary and sufficient for targeting Nav1.6 sodium channels to axon initial segments and nodes of Ranvier. *J. Neurosci.* 32:7232-43 (2012).
7. Rush AM, et al. Differential modulation of sodium channel Nav1.6 by two members of the fibroblast growth factor homologous factor 2 subfamily. *Eur. J. Neurosci.* 23:2551-62 (2006).
8. Eckenstein FP, McGovern T, Kern D, Deignan J. Neuronal vulnerability in transgenic mice expressing an inducible dominant-negative FGF receptor. *Exp. Neurol.* 198:338-49 (2006).

| Applications | Host | Isotype | Molecular Wt. | Species Cross-Reactivity |
|--------------|------|---------|--------------------|--------------------------|
| WB, IF/ICC | Goat | | 280kDa by SDS-PAGE | Hu, Rt, Ms |



Western blot analysis of whole brain lysates using goat pAb to MAP2, GPCA-MAP2, dilution 1:5,000 in green; [1] protein standard (red), [2] rat, and [3] mouse brain lysate. A band at about 280kDa corresponds to the MAP2A and MAP2B proteins.



Immunofluorescent analysis of rat brain stem section stained with goat pAb to MAP2, GPCA-MAP2, dilution 1:5,000 in red, and costained with mouse mAb to MBP, [MCA-7D2](#), dilution 1:5,000, in green. Following transcardial perfusion of rat with 4% paraformaldehyde, brain was post fixed for 24 hours, cut to 45μm, and free-floating sections were stained with above antibodies. The GPCA-MAP2 antibody labels MAP2 protein in the perikarya and dendrites of the most neurons, notably motoneurons in the brain stem, and the MBP antibody stains myelin sheath around axons.

Background:

Microtubules are 25nm diameter protein rods found in most kinds of eukaryotic cells and are associated with a family of proteins called microtubule associated proteins (MAPs). MAPs play a crucial role in the regulation of microtubule dynamics and interactions *in vivo*. MAP2 was originally named as one of the higher molecular weight MAPs with an SDS-PAGE molecular weight of about 280kDa (1-3). There is a single mammalian MAP2 gene which may generate two high molecular weight proteins of ~280kDa name MAP2A and MAP2B and multiple lower molecular weight forms usually named MAP2C and MAP2D which run on SDS-PAGE gels at 60-70kDa. The lower molecular weight forms are found in neurons early in development, but later are replaced by the higher molecular weight forms (2). The MAP2A and MAP2B forms include a protein insert missing in MAP2C and MAP2D which forms fine filamentous protrusions from the sides of brain microtubules referred to as the "projection domain". This antibody was made against the projection domain sequences and so is specific for MAP2A and MAP2B. MAP2 isoforms are expressed only in neuronal perikarya and dendrites so appropriate antibodies are used to identify neurons and dendrites in cell culture and sections (for example 4-8 used EnCor MAP2 antibodies).

The GPCA-MAP2 antibody was made against a mixture of recombinant human projection domain sequences, amino acids 377-1505, EnCor products [Prot-r-MAP2-P1](#), [Prot-r-MAP2-P2](#) and [Prot-r-MAP2-P3](#). It binds to the MAP2A and MAP2B isoforms but not the MAP2C and MAP2D which lack projection domain sequences. EnCor markets a mouse monoclonal antibody specific for sequence found in all MAP2 isoforms, [MCA-2C4](#), and also monoclonal antibodies binding epitopes only in MAP2A/B [MCA-4H5](#) and [MCA-5H11](#). We also market chicken and rabbit polyclonal antibodies to MAP2A/B with properties similar to this goat antibody, [CPCA-MAP2](#) and [RPCA-MAP2A/B](#).

FOR RESEARCH USE ONLY. NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE.

Abbreviation Key:

mAb—Monoclonal Antibody pAb—Polyclonal Antibody WB—Western Blot IF—Immunofluorescence ICC—Immunocytochemistry
 IHC—Immunohistochemistry E—ELISA Hu—Human Mo—Monkey Do—Dog Rt—Rat Ms—Mouse Co—Cow Pi—Pig Ho—Horse Ch—Chicken
 Dr—D. rerio Dm—D. melanogaster Sm—S. mutans Ce—C. elegans Sc—S. cerevisiae Sa—S. aureus Ec—E. coli.