


**Ordering Information**  
 Web [www.encorbio.com](http://www.encorbio.com)  
 Email [admin@encorbio.com](mailto:admin@encorbio.com)  
 Phone 352-372-7022  
 Fax 352-372-7066

**HGNC Name:** NEFL  
**UniProt:** P07196  
**RRID:** AB\_2861179  
**Immunogen:** C-terminal peptide of rat NF-L protein with an N-terminal Cys for coupling to KLH  
**Format:** Affinity purified at 1mg/mL in 50% PBS, 50% Glycerol plus 5mM Na3  
**Storage:** Store at 4°C, for longer term, store at -20°C  
**Recommended dilutions:**  
 WB: 1:10,000-1:20,000. IF/ICC: 1:5,000. IHC:1:5,000


**References:**

- Hoffman et al. Neurofilament gene expression: a major determinant of axonal caliber. *PNAS* 84:3472-6 (1987).
- Perrot R, et al. Review of the Multiple Aspects of Neurofilament Functions, and their Possible Contribution to Neurodegeneration. *Mol. Neurobiol.* 38:27-65 (2008).
- Lépinoux-Chambaud C, Eyer J. Review on intermediate filaments of the nervous system and their pathological alterations. *Histochem. Cell Biol.* 140:13-22 (2013).
- Liu Q, et al. Neurofilamentopathy in Neurodegenerative Diseases. *Open Neurol. J.* 5:58-62 (2011).
- Bacioglu M, et al. Neurofilament light chain in blood and CSF as marker of disease progression in mouse models and in neurodegenerative diseases. *Neuron* 91:56-66 (2016).

Applications	Host	Isotype	Molecular Wt.	Species Cross-Reactivity
WB, IF/ICC, IHC	Rabbit		68-70kDa	Hu, Rt, Ms, Co, Pi



Western blot analysis of tissue lysates using rabbit pAb to NF-L, RPCA-NF-L-ct, dilution 1:20,000, in green. [1] protein molecular weight standard (red), and lysates of rat brain [2], rat spinal cord [3], mouse brain [4] and mouse spinal cord [5]. The strong band at 68kDa corresponds to the NF-L protein.



Immunofluorescent analysis of rat cerebellum section stained with rabbit pAb to NF-L RPCA-NF-L-ct, dilution 1:5,000 in red, and costained with mouse mAb to beta-synuclein, MCA-6A10, dilution 1:500 in green. The blue is Hoechst staining of nuclear DNA. Following transcardial perfusion with 4% paraformaldehyde, the brain was post fixed for 24 hours, cut to 45µM, and free-floating sections were stained with above antibodies. The NF-L antibody labels dendrites and axons of neuronal cells, and beta-synuclein antibody detects protein that is concentrated in synaptic regions.

**Background:**

**Neurofilaments** are the 10nm or intermediate filament proteins found specifically in neurons, and are composed predominantly of three major proteins called NF-L, NF-M and NF-H, though other filament proteins may be included also. The major function of neurofilaments is likely to control the diameter of large axons (1). NF-L is the neurofilament light or low molecular weight polypeptide and runs on SDS-PAGE gels at 68-70kDa with some variability across species. Antibodies to NF-L are useful for identifying neuronal cells and their processes in cell culture and sectioned material. NF-L antibody can also be useful for the visualization of neurofilament rich accumulations seen in many neurological diseases, such as Lou Gehrig's disease (ALS), giant axon neuropathy, Charcot-Marie-Tooth disease and others (2-4). Much interest has recently been focused on the detection of NF-L released from neurons into blood and CSF as a surrogate marker of primarily axonal loss in a variety of types of CNS injury and degeneration (5).

RPCA-NF-L-ct was made against amino acids the C-terminal peptide, amino acids 515-543 of rat NF-L in NCBI entry [XP\\_032773476](https://www.ncbi.nlm.nih.gov/nuccore/XP_032773476), with an N-terminal cysteine added by which it was coupled to KLH. The antibody binds NF-L from a variety of species including human, rat and mouse and is useful for studies of neurofilament expression and proteolysis. In addition the antibody may be useful for ELISA studies as the epitopic region is known. We recently found that the epitope for this antibody is rapidly degraded during neurodegeneration, see our recent [BioRxiv](#) and [Brain Communications](#) papers for details. The antibody also works very well on paraffin embedded and formalin fixed human and rodent sections, see data under the "additional info" tag. We market several other NF-L antibodies including rabbit and chicken polyclonals [RPCA-NF-L](#) and [CPCA-NF-L](#), both made against full length recombinant human NF-L. We also have several mouse monoclonal antibodies to NF-L including epitope mapped [MCA-DA2](#), [MCA-6H63](#), [MCA-1B11](#) and [MCA-1D44](#).

**FOR RESEARCH USE ONLY. NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE.**

**Abbreviation Key:**

**mAb—Monoclonal Antibody pAb—Polyclonal Antibody WB—Western Blot IF—Immunofluorescence ICC—Immunocytochemistry IHC—Immunohistochemistry E—ELISA Hu—Human Mo—Monkey Do—Dog Rt—Rat Ms—Mouse Co—Cow Pi—Pig Ho—Horse Ch—Chicken Dr—D. rerio Dm—D. melanogaster Sm—S. mutans Ce—C. elegans Sc—S. cerevisiae Sa—S. aureus Ec—E. coli.**