

GFAP Chicken Polyclonal Antibody


CPCA-GFAP

Ordering Information
Web www.encorbio.com
Email admin@encorbio.com
Phone 352-372-7022
Fax 352-372-7066


HGNC Name: GFAP
UniProt: P14136
RRID: AB_2109953
Immunogen: Recombinant full length human GFAP isotype 1 expressed in and purified from *E. coli*.
Format: Concentrated IgY preparation in PBS plus 0.02% NaN₃
Storage: Store at 4°C
Recommended dilutions:
WB: 1:5,000. IF/ICC: 1:1,000-1:5,000 IHC: 1:20,000

References:
1. Bignami A, Eng LF, Dahl D, Uyeda CT. Localization of the glial fibrillary acidic protein in astrocytes by immunofluorescence. *Brain Res.* 43:429-35 (1972). 2. Yen SH, Fields KL. Antibodies to neurofilament, glial filament, and fibroblast intermediate filament proteins bind to different cell types of the nervous system. *J Cell Biol.* 88:115-26 (1981). 3. Shaw G, Osborn M, Weber K. An immunofluorescence microscopical study of the neurofilament triplet proteins, vimentin and glial fibrillary acidic protein within the adult rat brain. *Eur. J. Cell Biol.* 26:68-82 (1981). 4. Fitch MT, Silver J. CNS injury, glial scars, and inflammation: Inhibitory extracellular matrices and regeneration failure. *Exp. Neurol.* 209:294-301 (2008). 5. Brenner M, et al. Mutations in GFAP, encoding glial fibrillary acidic protein, are associated with Alexander disease. *Nat. Genet.* 27:117-20 (2001). The antibody has been sold through many OEM partners, and peer-reviewed publications making use of it can be found by searching Google Scholar for "CPCA-GFAP" or, if you are viewing this online, simply by selecting [this link](#).

Applications	Host	Isotype	Molecular Wt.	Species Cross-Reactivity
WB, IF/ICC, IHC	Chicken		~55kDa	Hu, Rt, Ms, Co, Pi, Ho



Western blot analysis of whole brain lysates using chicken pAb to GFAP, CPCA-GFAP, dilution 1:5,000 in green: [1] protein standard (red), [2] rat brain, [3] mouse brain. The strong band at about 50 kDa corresponds to the GFAP protein. Smaller proteolytic fragments and alternate transcripts of GFAP may also be detected on such blots.



Immunofluorescent analysis of a section of mouse hippocampus stained with chicken pAb to GFAP, CPCA-GFAP, dilution 1:5,000 in green and costained with rabbit pAb to FOX3/NeuN, RPCA-FOX3, dilution 1:5,000, in red. The blue is Hoechst staining of nuclear DNA. Following transcardial perfusion with 4% paraformaldehyde, mouse brain was post fixed for 24 hours, cut to 45μm, and free-floating sections were stained with the above antibodies. The GFAP antibody stains a network of astroglial cells while the Fox3/NeuN antibody stains the nuclei and proximal perikarya of neurons.

Background:

Glial Fibrillary Acidic Protein (GFAP) is a major CNS protein which runs on SDS-PAGE as a ~55kDa protein, usually associated with somewhat lower molecule weight bands which are alternate transcripts from the single gene. GFAP is strongly and specifically expressed in astrocytes and certain other glia in the central nervous system, in satellite cells in peripheral ganglia, and in non-myelinating Schwann cells in peripheral nerves and is also a component of neural stem cells (1-3). Astrocytes respond to many damage and disease states resulting in "astrogliosis" or the presence of a "glial response". GFAP antibodies are widely used to see the reactive astrocytes which form part of this response, since reactive astrocytes stain much more strongly with GFAP antibodies than normal astrocytes. GFAP also forms a major component of the so-called glial scar, an astrocyte rich structure apparently forming part of the barrier to nerve fiber regeneration following damage in the central nervous system (4). Neural stem cells frequently strongly express GFAP though they lose this if they develop into neurons or oligodendrocytes. Finally, Alexander disease was recently shown to be caused by point mutations in the protein coding region of the GFAP gene (5). All forms of Alexander disease are characterized by the presence of Rosenthal fibers, which are GFAP containing cytoplasmic inclusions found in astrocytes. Antibodies to GFAP are therefore very useful as a marker of glial cells in central and peripheral nerve system, as well as of developing neural stem cells. The CPCA-GFAP antibody was made against full length human recombinant GFAP, [Prot-r-GFAP](#), expressed in and purified from *E. coli*. The antibody works well on western blots, on immunostaining of cell culture or tissue sections and on paraffin embedded formalin fixed material. The same GFAP immunogen was used to produce rabbit, [RPCA-GFAP](#), and goat, [GPCA-GFAP](#) polyclonal antibodies. Using different immunogens, EnCor manufactures widely used mouse monoclonal antibodies to GFAP, [MCA-5C10](#), [MCA-2A5](#), and [MCA-3E10](#).

FOR RESEARCH USE ONLY. NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE.

Abbreviation Key:

mAb—Monoclonal Antibody pAb—Polyclonal Antibody WB—Western Blot IF—Immunofluorescence ICC—Immunocytochemistry IHC—Immunohistochemistry E—ELISA Hu—Human Mo—Monkey Do—Dog Rt—Rat Ms—Mouse Co—Cow Pi—Pig Ho—Horse Ch—Chicken Dr—D. rerio Dm—D. melanogaster Sm—S. mutans Ce—C. elegans Sc—S. cerevisiae Sa—S. aureus Ec—E. coli.