


Ordering Information
 Web www.encorbio.com
 Email admin@encorbio.com
 Phone 352-372-7022
 Fax 352-372-7066

HGNC Name: N.A.
UniProt: P0DTC2
RRID: AB_2861174
Immunogen: Recombinant SARS-CoV2 S-Protein ACE2 binding domain expressed in and purified from *E. coli*, EnCor product **PROT-SARS-CoV2-bd**
Format: Purified antibody at 1mg/mL in 50% PBS, 50% glycerol plus 5mM NaN₃
Storage: Shipped on ice. Store at 4°C for short term, for longer term at -20°C. Avoid freeze / thaw cycles.
Recommended dilutions:
 WB: 1:1,000-1:3,000. ICC/IF: 1:1,000


References:

1. Wu, F et al. A new coronavirus associated with human respiratory disease in China. *Nature* doi:10.1038/s41586-020-2008-3-2020 579:265-269 (2020).
2. Ren, L-L et al. Identification of a novel coronavirus causing severe pneumonia in human: a descriptive study. *Chin Med J (Engl)* doi:10.1097/CM9.0000000000000722 133:1015-24 (2020).
3. Walls, A C et al. Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell* doi:10.1016/j.cell.2020.02.058 180:1-12 (2020)
4. Yan, R et al. Structural basis for the recognition of SARS-CoV-2 by full-length human ACE2. *Science* doi:10.1126/science.abb2762 367:1444-8 (2020).
5. Wang, D-S et al. The pleckstrin homology domain of human beta I sigma II spectrin is targeted to the plasma membrane in vivo. *Biochem. Biophys. Res. Comm.* 225:420-6 (1996).

Applications	Host	Isotype	Molecular Wt.	Species Cross-Reactivity
WB, ICC/IF, IHC	Mouse	IgG1	Intact S-Protein 142kDa	NA



Left Panel: Western blot analysis of HEK293 cell lysates using mouse mAb to SARS-CoV2-bd protein, MCA-5G8, dilution 1:3,000 in green: [1] protein standard, [2] non-transfected cells, [3] cells transfected with pCI-Neo-Mod containing the SARS-CoV2-bd cDNA, and [4] cells transfected with pCI-Neo-GFP vector expression construct containing the SARS-CoV2-bd cDNA. The band at 25kDa mark in the transfected cells demonstrates expression of SARS-CoV-bd protein, and the band at about 50kDa corresponds to a GFP-SARS-CoV2-bd fusion protein. The same blot was simultaneously probed with EnCor rabbit pAb to HSP60, **RPCA-HSP60**, dilution 1:5,000, in red, revealing a single band at 60kDa in both transfected and non-transfected cells. Right Panel: Blot of full length recombinant SARS-CoV2 S-protein expressed in HEK293 cells, product **10561-CV**, obtained from **R&D Systems**. Lane 6 shows a loading of 1µg and lane 7 is 100ng. On longer exposure of the blot the antibody could readily detect 10ng of the S-protein. Lanes 1 and 5 are molecular weight standards of indicated size.



Immunofluorescent analysis of HEK293 cells transfected with pCI-Neo-Mod vector (5) including SARS-CoV2-bd cDNA and stained with mouse mAb to SARS-CoV2-bd, MCA-5G8, dilution 1:1,000, in red. The blue is Hoechst staining of nuclear DNA. The MCA-5G8 antibody reveals expression of SARS-CoV2-bd protein only in transfected cells. DAPI reveals the nuclear DNA of both transfected and non-transfected cells.

Background:

In late 2019 a novel infectious disease was discovered in Wuhan, China which was quickly recognized to be caused by a previously unknown RNA coronavirus. The virus was very rapidly isolated, the full RNA sequence determined and put on-line on the 10th of January 2020. The sequence revealed that the virus was most closely related to certain bat coronaviruses and the severe acute respiratory syndrome (SARS) coronavirus. Immediately biotechnology companies and research institutes used the RNA sequence information to generate vaccine candidates. The SARS virus was known to enter and infect human cells by means of the so-called spike or S-protein which binds to the extracellular domain of the angiotensin converting enzyme 2 (ACE2) protein, which is then internalized bringing the virus into the cell. Cryoelectron microscopy and binding studies quickly determined that the S-protein of SARS-CoV2 is structurally similar to that of the SARS virus and also binds to the ACE2 receptor, albeit with higher affinity than the S-protein of SARS. This focuses attention on the ACE2 binding site on the SARS-CoV2 S-protein and for the complementary region on ACE2 which binds the SARS-CoV2 S-protein. We therefore expressed both these regions in *E. coli*, our products **PROT-SARS-CoV2-bd** and **PROT-ACE2-bd** and raised antibodies to them.

The MCA-5G8 antibody was made against our recombinant construct comprising amino acids 308-541 in the S-protein sequence in **SARS-CoV2 Wuhan-Hu-1, complete genome**. The antibody works well on western blots of crude homogenates of HEK293 cells transfected with the SARS-CoV2 binding domain, cleanly producing the appropriate sized band and as expected also binds the full length S-protein. In addition S-protein transfected cells and cells infected with patient derived SARS-CoV2 show clean and strong immunofluorescence staining of transfected or infected cells. We are currently determining the exact peptide epitope of this and our other SARS-CoV2 S-protein antibodies and also measuring their kinetic properties. EnCor supplies another mouse monoclonal antibody to the SARS-CoV2 S-protein ACE2 binding domain **MCA-2G1** and also a rabbit polyclonal **RPCA-SARS-CoV2-bd**.

FOR RESEARCH USE ONLY. NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE.

Abbreviation Key:

mAb—Monoclonal Antibody pAb—Polyclonal Antibody WB—Western Blot IF—Immunofluorescence ICC—Immunocytochemistry IHC—Immunohistochemistry E—ELISA Hu—Human Mo—Monkey Do—Dog Rt—Rat Ms—Mouse Co—Cow Pi—Pig Ho—Horse Ch—Chicken Dr—D. rerio Dm—D. melanogaster Sm—S. mutans Ce—C. elegans Sc—S. cerevisiae Sa—S. aureus Ec—E. coli.

FOR RESEARCH USE ONLY. NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE.

Abbreviation Key:

mAb—Monoclonal Antibody *pAb*—Polyclonal Antibody *WB*—Western Blot *IF*—Immunofluorescence *ICC*—Immunocytochemistry
IHC—Immunohistochemistry *E*—ELISA *Hu*—Human *Mo*—Monkey *Do*—Dog *Rt*—Rat *Ms*—Mouse *Co*—Cow *Pi*—Pig *Ho*—Horse *Ch*—Chicken
Dr—*D. rerio* *Dm*—*D. melanogaster* *Sm*—*S. mutans* *Ce*—*C. elegans* *Sc*—*S. cerevisiae* *Sa*—*S. aureus* *Ec*—*E. coli*.