

SARS-CoV2 S-protein ACE2 binding domain


Mouse Monoclonal Antibody

MCA-2G1

Ordering Information
 Web www.encorbio.com
 Email admin@encorbio.com
 Phone 352-372-7022
 Fax 352-372-7066

HGNC Name: NA
UniProt: P0DTC2
RRID: AB_2861173
Immunogen: Recombinant SARS-CoV2 S-Protein ACE2 binding domain expressed in and purified from *E. coli*, EnCor product PROT-r-SARS-CoV2-bd
Format: Purified antibody at 1mg/mL in 50% PBS, 50% glycerol plus 5mM Na₂S₂O₅
Storage: Shipped on ice. Store at 4°C for short term, for longer term at -20°C. Avoid freeze / thaw cycles.
Recommended dilutions:
 WB: 1:1,000-1:3,000. ICC/IF: 1:1,000

References:
 1. Wu, F et al. A new coronavirus associated with human respiratory disease in China. *Nature* doi:10.1038/s41586-020-2008-3.2020 579:265-269 (2020).
 2. Ren, L-L et al. Identification of a novel coronavirus causing severe pneumonia in human: a descriptive study. *Chin Med J (Engl)* doi:10.1097/CM9.0000000000000722 133:1015-24 (2020).
 3. Walls, A C et al. Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell* doi:10.1016/j.cell.2020.02.058 180:1-12 (2020)
 4. Yan, R et al. Structural basis for the recognition of SARS-CoV-2 by full-length human ACE2. *Science* doi:10.1126/science.abb2762 367:1444-8 (2020).
 5. Wang, D-S et al. The pleckstrin homology domain of human β -I α -II spectrin is targeted to the plasma membrane in vivo. *Biochem. Biophys. Res. Comm.* 225:420-6 (1996).



Left Panel: HEK293 cells were transfected with DNA encoding the S-protein ACE2 binding site in **PROT-SARS-CoV2-bd** which was inserted into pCI-Neo-Mod or pCI-Neo-GFP eukaryotic expression vectors, which express either the insert only or the insert fused with GFP (5). Lane 2 shows a crude homogenate of untransfected control cells, lane 3 shows a homogenate of cells expressing SARS-CoV2-bd and lane 4 shows a homogenate of cells expressing GFP-SARS-CoV2-bd fusion. The MCA-2G1 was used at a dilution of 1:3,000 and produces clean and specific staining of bands of the expected size as shown in green. The blot was also stained with EnCor rabbit polyclonal control antibody to HSP60, **RPCA-HSP60**, at a dilution of 1:5,000 in red. Right Panel: Blot of full length recombinant SARS-CoV2 S-protein expressed in HEK293 cells, product **10561-CV**, obtained from **R&D Systems**. Lane 6 shows a loading of 1 μ g and lane 7 is 100ng. On longer exposure of the blot the antibody could readily detect 10ng of the S-protein. Lanes 1 and 5 are molecular weight standards of indicated size.

HEK293 cells were transfected with the DNA encoding the S-protein segment in **PROT-SARS-CoV2-bd** which was inserted in the pCI-Neo-Mod expression vector (5). The MCA-2G1 produces clean and specific staining of transfected cells which stain identically with **RPCA-SARS-CoV2-bd**, our rabbit polyclonal antibody to the same immunogen. The nuclei of transfected and untransfected cells are shown in blue with DAPI DNA stain.

Background:

In late 2019 a novel infectious disease was discovered in Wuhan, China which was quickly recognized to be caused by a previously unknown RNA coronavirus. The virus was very rapidly isolated, the full RNA sequence determined and put on-line on the 10th of January 2020. The sequence revealed that the virus was most closely related to certain bat coronaviruses and the severe acute respiratory syndrome (SARS) coronavirus. Immediately biotechnology companies and research institutes used the RNA sequence information to generate vaccine candidates. The SARS virus was known to enter and infect human cells by means of the so-called spike or S-protein which binds to the extracellular domain of the angiotensin converting enzyme 2 (ACE2) protein, which is then internalized bringing the virus into the cell. Cryoelectron microscopy and binding studies quickly determined that the S-protein of SARS-CoV2 is structurally similar to that of the SARS virus and also binds to the ACE2 receptor, albeit with higher affinity than the S-protein of SARS. This focuses attention on the ACE2 binding site on the SARS-CoV2 S-protein and for the complementary region on ACE2 which binds the SARS-CoV2 S-protein. We therefore expressed both these regions in *E. coli*, our products **PROT-R-SARS-CoV2-bd** and **PROT-R-ACE2-bd** and raised antibodies to them.

The MCA-2G1 antibody was made against our recombinant construct comprising amino acids 308-541 in the S-protein sequence in **SARS-CoV2 Wuhan-Hu-1, complete genome**. The antibody works well on western blots of crude homogenates of HEK293 cells transfected with the SARS-CoV2 binding domain, cleanly producing the appropriate sized band and as expected also binds the full length S-protein. In addition S-protein transfected cells show clean and strong immunofluorescence staining of the expressed protein with this antibody. We are currently determining the exact peptide epitope of this and our other SARS-CoV2 S-protein antibodies and also measuring their kinetic properties. EnCor supplies another mouse monoclonal antibody to the SARS-CoV2 S-protein ACE2 binding domain **MCA-5G8** and also a rabbit polyclonal **RPCA-SARS-CoV2-bd**.

FOR RESEARCH USE ONLY. NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE.

Abbreviation Key:

mAb—Monoclonal Antibody **pAb**—Polyclonal Antibody **WB**—Western Blot **IF**—Immunofluorescence **ICC**—Immunocytochemistry
IHC—Immunohistochemistry **E**—ELISA **Hu**—Human **Mo**—Monkey **Do**—Dog **Rt**—Rat **Ms**—Mouse **Co**—Cow **Pi**—Pig **Ho**—Horse **Ch**—Chicken
Dr—*D. rerio* **Dm**—*D. melanogaster* **Sm**—*S. mutans* **Ce**—*C. elegans* **Sc**—*S. cerevisiae* **Sa**—*S. aureus* **Ec**—*E. coli*.

FOR RESEARCH USE ONLY. NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE.

Abbreviation Key:

*mAb—Monoclonal Antibody pAb—Polyclonal Antibody WB—Western Blot IF—Immunofluorescence ICC—Immunocytochemistry
IHC—Immunohistochemistry E—ELISA Hu—Human Mo—Monkey Do—Dog Rt—Rat Ms—Mouse Co—Cow Pi—Pig Ho—Horse Ch—Chicken
Dr—D. rerio Dm—D. melanogaster Sm—S. mutans Ce—C. elegans Sc—S. cerevisiae Sa—S. aureus Ec—E. coli.*