


Ordering Information
Web www.encorbio.com
Email admin@encorbio.com
Phone 352-372-7022
Fax 352-372-7066

HGNC Name: NA
UniProt: NA
RRID: AB_2889159
Immunogen: Recombinant CasΦ/Cas-12j expressed in and purified from *E. coli*
Format: Purified antibody at 1mg/mL in 50% PBS, 50% glycerol plus 5mM Na₂S₂O₃
Storage: Shipped on ice. Store at 4°C for short term, for longer term at -20°C. Avoid freeze / thaw cycles.
Recommended dilutions:
WB: 1:1,000-1:2,000. ICC/IF: 1:1,000

References:

1. Al-Shayeb, B et al. Clades of huge phages from across Earth's ecosystems. *Nature* DOI: 10.1038/s41586-020-2007-4 578:425-531 (2020).
2. Pausch, P. CRISPR-CasΦ from huge phages is a hypercompact genome editor. *Science* DOI: 10.1126/science.abb1400 369:333-337 (2020).



Background:

There has been much recent interest in gene editing and other genetic manipulations by CRISPR-Cas family enzymes. Recent ecosystem and microbiome DNA sequencing and assembly characterized genomes of "huge phages" or megaphages, see Al-Shayeb et al. *Nature* 578:425-431 (2020) and showed that they contained novel Cas family enzymes. The lab of Nobel prize winner Jennifer Doudna characterized these Cas enzymes and showed that they could be used to perform CRISPR, but had the significant advantage that they were much smaller in molecular size than other Cas family enzymes such as the widely used Cas9 enzymes from *S. pyogenes* and *S. aureus*, see Pausch, P. et al. *Science* 369:333-337 (2020). One of these enzymes is referred to as CasΦ and also known as Cas-12j and was the focus of the Pausch et al. paper. The use of such smaller Cas family enzymes leaves more room for other nucleic acid sequences in AAV and other viral vectors which typically have a limited DNA capacity, thus allowing more versatility for new CRISPR based manipulations.

The MCA-5F95 antibody was made against our recombinant construct of the full sequence of CasΦ-2 from Pausch et al. 2020. The antibody works well on western blots of crude homogenates of HEK293 cells transfected with the Cas-Φ-2 DNA, cleanly producing the appropriate sized band. In addition such transfected cells show clean and strong cytoplasmic staining by immunofluorescence staining with this antibody. We also supply a rabbit polyclonal antibody to this protein, [RPCA-Cas12j](#).

FOR RESEARCH USE ONLY. NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE.

Abbreviation Key:

mAb—Monoclonal Antibody pAb—Polyclonal Antibody WB—Western Blot IF—Immunofluorescence ICC—Immunocytochemistry
IHC—Immunohistochemistry E—ELISA Hu—Human Mo—Monkey Do—Dog Rt—Rat Ms—Mouse Co—Cow Pi—Pig Ho—Horse Ch—Chicken
Dr—D. rerio Dm—D. melanogaster Sm—S. mutans Ce—C. elegans Sc—S. cerevisiae Sa—S. aureus Ec—E. coli.